

## MODELING OF ELECTRON DYNAMICS IN DNA

**V.D. Lakhno**

*Institute of Mathematical Problems of Biology, Russian Academy of Sciences  
142290, Pushchino, Moscow Region, Russia  
lak@impb.psn.ru*

General properties of electron motion in molecular chains are considered. The charge motion is described in terms of quantum mechanics, whereas vibrational degrees-of-freedom are treated both classically and quantum mechanically. A typical charge transfer/transport pattern can physically be viewed as a polaron and/or soliton. A closed analytical expression for charge carrier velocity dependence on electric field has been derived and analyzed in detail. Special attention is given to: dynamical behavior of electrons in rigid chains, band structure of regular polynucleotide chains, dynamics of polaron states formation in Holstein chain, polaron motion in an electric field, the role of dispersion, Bloch oscillations and breather states, the bipolaron mechanism of a superconducting state in DNA based on Holstein-Hubbard model.

- [1] V.D.Lakhno, V.B.Sultanov, Bipolarons in homogeneous DNA duplexes, *Chem. Phys. Lett.*, v.503, (2011), pp.292-295.
- [2] V.D.Lakhno, A.N.Korshunova, Electron motion in a Holstein molecular chain in an electric field, *EPJB*, v.79, (2011), pp.147-151.
- [3] V.D.Lakhno, Davydov's Solitons in a Homogeneous Nucleotide Chain, *Int. Journ. Quant. Chem.*, v.110, (2010), pp.127-137.
- [4] V.D.Lakhno, Energy and Critical Ionic-Bond Parameter of a 3D-Large-Radius Bipolaron, *JETP*, v.137, (2010), pp.926-931.
- [5] N.I.Kashirina, V.D.Lakhno, Large-radius bipolaron and the polaron-polaron interaction, *Physics-Uspekhi*, v.53, (2010), pp.431-453.